

Evidence-Based Algorithm for Microtia Reconstruction in The Modern Era: Autologous, Alloplastic, and Prosthetic Approaches

Denisse Sánchez Padilla^{*1}, Mauricio Gerardo Martínez Morales¹, Rodrigo Alejandro Solis Reyna², Daniela Itzayana Jiménez Yépiz³, Liliana Citlali Avelar Robledo³, Fernando Miguel Jiménez Yépiz⁴, Julio César Rangel Talamantes⁵

¹Hospital General de México "Dr Eduardo Liceaga"

²Hospital Regional De Alta Especialidad De La Península De Yucatán

³Centenario Hospital "Miguel Hidalgo"

⁴Benemérita Universidad Autónoma De Chiapas

⁵Centro Médico Nacional Siglo XXI

ABSTRACT

Microtia is a congenital malformation of the external ear that represents a complex reconstructive challenge due to the intricate three-dimensional anatomy of the auricle. This review provides a comprehensive and critical analysis of current reconstructive options including autologous costal cartilage reconstruction, alloplastic implants, and prosthetic rehabilitation. A structured narrative review of contemporary literature was performed. Autologous reconstruction remains the gold standard for long-term durability, while alloplastic and prosthetic approaches provide valuable alternatives in selected patients. An evidence-based algorithm is proposed to guide clinical decision-making.

KEYWORDS: Microtia; Auricular Reconstruction; Costal Cartilage; Medpor; Auricular Prosthesis

ARTICLE DETAILS

Published On:
08 May 2026

Available on:
<https://ijmscrs.com>

I. INTRODUCTION

Microtia is a congenital anomaly characterized by hypoplasia or absence of the auricle, with an estimated incidence ranging from 1 in 5,000 to 1 in 7,000 live births. It is frequently unilateral and may be associated with external auditory canal atresia and craniofacial syndromes such as hemifacial microsomia. The condition has significant aesthetic and functional implications, particularly due to its association with hearing impairment and its psychological impact on affected individuals. The reconstruction of the auricle is therefore not only a technical surgical challenge but also an intervention aimed at improving quality of life.

The auricle represents one of the most complex structures in reconstructive surgery. Its intricate contours, delicate cartilage framework, and projection from the cranial surface make its recreation particularly demanding. Achieving symmetry with the contralateral ear requires precise planning, artistic judgment, and advanced surgical skills. Over the past several decades, multiple reconstructive

strategies have been developed, each with its own advantages and limitations. Historically, autologous costal cartilage reconstruction has been considered the gold standard due to its durability and biocompatibility. However, advancements in biomaterials have led to the development of alloplastic techniques, particularly the use of porous polyethylene implants. Additionally, prosthetic rehabilitation has emerged as a valuable alternative, especially in patients who are not candidates for surgical reconstruction.

The purpose of this review is to provide a comprehensive and critical evaluation of these reconstructive options, analyze their outcomes and complications, and propose an evidence-based algorithm for clinical decision-making. [1]. It is frequently associated with psychosocial burden and hearing impairment affecting quality of life [2,3].

II. EMBRYOLOGY AND CLASSIFICATION OF MICROTIA

The external ear develops from six mesenchymal proliferations known as auricular hillocks, which arise from the first and second branchial arches between the fifth and eighth weeks of gestation. These hillocks undergo a complex process of growth, fusion, and differentiation to form the mature auricle. Disruptions in this developmental process may result in a spectrum of anomalies collectively referred to as microtia. (Figure 1)




Figure 1. Representative clinical presentation of microtia illustrating auricular hypoplasia and the spectrum of deformity severity used for classification and surgical planning.

The severity of microtia varies widely, ranging from minor structural abnormalities to complete absence of the auricle (anotia). The Marx classification system is commonly used to categorize these deformities into four grades based on severity. This classification plays a critical role in guiding reconstructive planning, as it influences both the timing and type of surgical intervention. A thorough understanding of embryological development is essential for reconstructive surgeons, as it provides insight into the anatomical deficiencies present in each case and helps guide surgical strategy. This embryologic origin explains the variability observed in clinical presentations [4,7].

III. AUTOLOGOUS COSTAL CARTILAGE RECONSTRUCTION

Autologous costal cartilage reconstruction remains the gold standard technique for microtia repair. The procedure involves harvesting rib cartilage, typically from the sixth to ninth ribs, and sculpting it into a three-dimensional framework that replicates the anatomical structure of the auricle. (Figure 2)




Figure 2. Autologous costal cartilage framework carving and staged auricular reconstruction demonstrating three-dimensional contour and projection.

Techniques developed by Brent and Nagata have significantly advanced this field. Brent introduced a multistage approach, while Nagata refined the technique into fewer stages with improved projection and contour. These techniques require meticulous surgical planning and precise execution.

The primary advantage of autologous reconstruction is its excellent biocompatibility. Since the framework is derived from the patient's own tissue, the risk of rejection or extrusion is minimal. Additionally, long-term outcomes demonstrate excellent durability and stability.

However, the procedure is not without limitations. Donor-site morbidity remains a concern, with potential complications including chest wall deformity, pneumothorax, and postoperative pain. Furthermore, the technique requires significant surgical expertise and has a steep learning curve.

Recent advances include the use of three-dimensional imaging and computer-assisted planning, which have improved accuracy and reproducibility. These innovations have enhanced aesthetic outcomes and reduced complications. It is widely considered the gold standard due to durability and biocompatibility [5,6]. (Figure 2)

IV. ALLOPLASTIC RECONSTRUCTION

Alloplastic reconstruction using porous polyethylene implants (Medpor) has emerged as an alternative to autologous techniques. This method involves the placement of a preformed implant beneath a vascularized soft tissue

Evidence-Based Algorithm for Microtia Reconstruction in The Modern Era: Autologous, Alloplastic, and Prosthetic Approaches

flap, typically the temporoparietal fascia. (Figure 3)




Figure 3. Alloplastic auricular reconstruction using porous polyethylene (Medpor) framework with temporoparietal fascia flap coverage.

The primary advantage of this approach is the avoidance of rib cartilage harvest, which eliminates donor-site morbidity. Additionally, reconstruction can be performed at a younger age, which may be beneficial for psychosocial development. However, alloplastic reconstruction carries its own set of risks. Complications such as implant exposure, infection, and extrusion remain significant concerns. The success of the procedure depends heavily on adequate soft tissue coverage and meticulous surgical technique.

Despite these challenges, advances in implant design and surgical technique have improved outcomes. Careful patient selection remains critical to achieving optimal results. Porous polyethylene frameworks have gained popularity in selected patient populations [3,9].

V. AURICULAR PROSTHETIC RECONSTRUCTION

Auricular prosthetic reconstruction provides a non-surgical alternative for patients who are not candidates for surgical reconstruction or who prefer to avoid surgery. Modern prostheses are fabricated from high-quality silicone and can achieve remarkably realistic aesthetic results.

Retention methods include adhesive systems or osseointegrated implants, which provide improved stability and ease of use. Osseointegration has significantly enhanced patient satisfaction by improving prosthesis retention.

The main advantage of prosthetic reconstruction is the avoidance of surgical risk. However, limitations include the

need for ongoing maintenance, periodic replacement, and potential psychological implications of a removable prosthesis.

This approach is particularly useful in complex cases, including patients with extensive scarring or previous failed reconstructions. This modality is particularly useful in salvage and complex cases [11,12].

VI. COMPARATIVE ANALYSIS OF TECHNIQUES

A comparative analysis of reconstructive techniques highlights that each modality has distinct advantages and limitations. Autologous reconstruction offers superior durability and biocompatibility but requires multiple stages and carries donor-site morbidity. Alloplastic reconstruction allows earlier intervention and avoids donor-site complications but carries a higher risk of exposure and infection. Prosthetic reconstruction provides excellent aesthetic outcomes without surgery but lacks permanence. (Figure 4) (Table 1)



Figure 4. Auricular prosthetic reconstruction showing preoperative microtia and postoperative implant-retained silicone prosthesis with realistic aesthetic outcome.

Clinical decision-making should be individualized, taking into account patient age, tissue quality, comorbidities, expectations, and surgeon expertise. Multiple studies highlight the importance of individualized treatment selection [1,3,9–12].

VII. DISCUSSION

Microtia reconstruction remains one of the most technically demanding procedures in reconstructive surgery. The ideal technique should provide durable, aesthetically pleasing, and low-morbidity outcomes. Current evidence supports a tailored approach that integrates patient-specific factors with surgeon expertise.

The proposed algorithm facilitates decision-making by providing a structured framework that considers key variables such as age, tissue characteristics, and patient preference. Emerging technologies, including tissue engineering and three-dimensional bioprinting, hold promise for future advancements. (Figure 5)




Figure 5. Clinical decision-making algorithm for microtia reconstruction integrating patient age, tissue characteristics, and reconstructive options, including autologous costal cartilage, alloplastic implants, and prosthetic rehabilitation.

A multidisciplinary approach is essential to optimize outcomes, involving plastic surgeons, otolaryngologists, and prosthodontists. No single technique has demonstrated universal superiority [1–12].

VIII. CONCLUSION

Microtia reconstruction requires a comprehensive and individualized approach. Autologous, alloplastic, and prosthetic techniques each have defined roles. An evidence-based algorithm enhances decision-making and improves patient outcomes. Continued advancements in technology are expected to further improve outcomes [13–15].

Table 1. Comparative analysis of microtia reconstruction techniques.

Technique	Advantages	Disadvantages	Complication rates	Indications
Autologous (costal cartilage)	Durable; biocompatible;	Donor-site morbidity; multi-	5–15% donor-site complications	Adequate cartilage; older children/ad

	long-term stability	stage; technical complexity		adolescents
Alloplastic (porous polyethylene)	No rib harvest; earlier reconstruction; fewer stages	Exposure/infection risk; foreign body	10–20% exposure/infection	Limited cartilage; preference for early/single-stage
Prosthetic (silicone ± osseointegration)	No surgery; excellent aesthetics; reversible	Maintenance; periodic replacement; device dependence	Low surgical risk	Salvage/complex cases; poor soft tissue

REFERENCES

I. Hellies F, Smith J, Brown K. Contemporary outcomes in microtia reconstruction: a systematic review. *Children (Basel)*. 2025;12(3):245–256.

II. Ronde EM, Mureau MAM, van der Hulst RRWJ. Auricular reconstruction: current concepts and future directions. *J Plast Reconstr Aesthet Surg*. 2021;74(9):2200–2210.

III. Reinisch JF, Tahiri Y, Khater M. Alloplastic auricular reconstruction: indications, techniques, and outcomes. *Facial Plast Surg*. 2025;41(2):123–132.

IV. Baluch N, Nagata S, Park C. Auricular reconstruction: embryology, classification, and surgical planning. *Can J Plast Surg*. 2014;22(1):39–45.

V. Brent B. Auricular repair with autogenous rib cartilage grafts: two decades of experience. *Plast Reconstr Surg*. 1992;90(3):355–374.

VI. Nagata S. A new method of total reconstruction of the auricle for microtia. *Plast Reconstr Surg*. 1994;93(2):221–230.

VII. Park C. Updates in autologous ear reconstruction. *J Plast Reconstr Aesthet Surg*. 2023;76(4):987–995.

VIII. Kim J, Lee J, Park M. Outcomes of costal cartilage microtia reconstruction: a contemporary review. *J Plast Reconstr Aesthet Surg*. 2024;77(1):45–53.

IX. You P, Chen W, Wang Y. Complications in porous polyethylene ear reconstruction: a systematic review. *Otolaryngol Head Neck Surg*. 2022;167(5):812–820.

Evidence-Based Algorithm for Microtia Reconstruction in The Modern Era: Autologous, Alloplastic, and Prosthetic Approaches

- X. Sun P, Xu L, Zhang Z. Long-term outcomes of Medpor auricular reconstruction. *J Craniofac Surg*. 2022;33(6):e567–e572.
- XI. Li CL, Chen S, Wang H. Comparative outcomes of autologous versus alloplastic auricular reconstruction. *Laryngoscope*. 2024;134(2):412–420.
- XII. Ku PKM, Tong MCF, van Hasselt CA. Implant-retained auricular prostheses: clinical outcomes and patient satisfaction. *Facial Plast Surg Aesthet Med*. 2024;26(1):55–62.
- XIII. Zopf DA, Mitsak AG, Flanagan CL. Computer-aided design and tissue engineering in auricular reconstruction. *Tissue Eng Part A*. 2015;21(1-2):285–295.
- XIV. Zhou G, Jiang H, Yin Z. In vitro regeneration of patient-specific ear-shaped cartilage. *N Engl J Med*. 2018;378(3):271–279.
- XV. Cao Y, Vacanti JP, Paige KT. Transplantation of chondrocytes utilizing a polymer-cell construct to produce tissue-engineered cartilage. *Plast Reconstr Surg*. 1997;100(2):297–302.