

Aesthetic Blepharoplasty and Periorbital Rejuvenation: A Comprehensive Review of Surgical Techniques, Risk Prevention, And Complication Management

Sergio Morales Acosta^{*1}, Sarahí Carrillo Barraza², María Fernanda Vázquez Páez³, Claudia Lizama Rubio⁴, Daniel Antonio Beltran Valdez⁵, María de Jesús Aphil Montiel Herrera⁶, Irvint Joel Bautista Perez⁷, Osiel González Villegas⁸

¹Centro Universitario de Ciencias de la Salud (CUCS).

²Hospital Universitario Torreón

³Hospital de Especialidades CMN Puebla.

⁴Hospital General “Fernando Quiroz Gutiérrez” ISSSTE, CDMX.

⁵Hospital General Fray Junipero Serra ISSSTE, Tijuana, Baja California.

⁶Hospital General de Atizapán “Dr. Salvador González Herrejon “, Atizapán de Zaragoza, Estado de México.

⁷Hospital general “Dr. Fernando Quiroz Gutiérrez” ISSSTE, CDMX.

⁸Universidad Cuauhtémoc, campus San Luis Potosí.

ABSTRACT

Contemporary literature on upper and lower aesthetic blepharoplasty, evaluating the evolution of surgical techniques, anatomical considerations, and strategies for complication management.

Methods: A literature review was conducted based on selected articles regarding surgical techniques (transconjunctival approaches, pinch technique, tarsal fixation), preoperative evaluation, and management of eyelid retraction. Studies focusing on tissue preservation and risk reduction were prioritized.

Results: The current surgical paradigm has transitioned from aggressive tissue resection to the preservation and repositioning of fat and the orbicularis oculi muscle. In upper blepharoplasty, the emphasis is on avoiding the "operated look" through precise incisions and volume preservation. In lower blepharoplasty, the transconjunctival approach combined with fat repositioning and lateral canthopexy proves superior in preventing eyelid malposition. The management of post-blepharoplasty lower eyelid retraction (PBLER) requires complex reconstructive approaches that avoid conventional skin grafts.

KEYWORDS: r eyelid, eyelid retraction, transconjunctival approach

ARTICLE DETAILS

Published On:
04 May 2026

Available on:
<https://ijmscrs.com>

I. INTRODUCTION

Aesthetic blepharoplasty has undergone a significant philosophical and technical evolution over the past three decades. Traditionally conceived as a subtractive procedure focused on the aggressive excision of skin, orbicularis muscle, and orbital fat, the technique often resulted in periorbital skeletonization that paradoxically aged the patient or caused long-term functional complications. The contemporary paradigm recognizes that periorbital youthfulness is associated with the maintenance of facial volume and a smooth transition at the lid-cheek junction.

Currently, blepharoplasty is among the most frequently performed facial cosmetic procedures globally, for both aesthetic and functional indications (e.g., severe dermatochalasis or fat pseudoherniation). This review aims to comprehensively analyze modern surgical approaches for upper and lower blepharoplasty, preoperative risk stratification, and the advanced management of challenging complications, such as lower eyelid retraction.

This document is a template. An electronic copy can be downloaded from the conference website. For questions on paper guidelines, please contact the conference publications

Aesthetic Blepharoplasty and Periorbital Rejuvenation: A Comprehensive Review of Surgical Techniques, Risk Prevention, And Complication Management

committee as indicated on the conference website. Information about final paper submission is available from the conference website. Achieving an ideal aesthetic outcome in blepharoplasty requires a deep understanding of the upper eyelid's surface landmarks. The lid crease is formed by the attachment of the levator aponeurosis to the subdermal tissues. Positioned just above this is the eyelid fold, which typically obscures the crease in a youthful eye. The area between the lid margin and the crease is known as the tarsal platform. Aesthetic harmony is often measured by the Golden Ratio (1:1.618), representing the ideal proportion between the tarsal platform and the distance from the crease to the eyebrow. Surgeons must be cautious: over-resecting fat can collapse this ratio toward 1:1, leading to a hollowed, aged look. Preserving the 3D structure of the lid fold is equally vital; excessive skin or fat removal can "flatten" the eye's profile, creating an artificial appearance.

Pre-Surgical Assessment: The Brow and Skin Quality

Temporal hooding is frequently a dual issue caused by lateral dermatochalasis and temporal brow ptosis. It is critical to distinguish if the patient requires a brow lift in conjunction with—or instead of—direct lid surgery.

- **Vertical Dimensions:** For a natural look, the distance from the lower brow border to the lid margin should ideally measure between 25 mm and 30 mm, and never less than 20 mm.
- **Skin Condition:** While eyelid skin (approx. 850 microns thick) does not thin significantly with age, it loses elasticity. While pretarsal skin is smooth in younger patients, elastin loss leads to the development of fine lines.
- **Systemic Factors:** Pronounced dermatochalasis may be exacerbated by conditions such as thyroid eye disease, Ehlers-Danlos syndrome, or chronic edema.

Internal Lamellar Structure

The eyelid is organized into functional layers or "lamellae," which vary depending on the vertical level:

1. **At the Tarsal Level:**
 - Anterior: Skin and orbicularis oculi muscle.
 - Posterior: Tarsal plate and conjunctiva.
2. **Above the Tarsal Plate:**
 - Anterior: Skin and orbicularis muscle.
 - Middle: The orbital septum.
 - Posterior: Levator aponeurosis, Müller's muscle, and conjunctiva.

Key Surgical Landmarks

The Orbital Septum and Fat Pads

The orbital septum serves as a vital protective barrier for the levator muscle. In Asian eyelids, the septum typically inserts lower on the aponeurosis.

Behind the septum lie two distinct fat pads:

- **Medial Fat Pad:** Usually paler and firmer.

- **Central Fat Pad:** More yellow (due to higher lutein/beta-carotene content) and more vascularized.

Musculature and Innervation

- **Levator Aponeurosis:** Under the influence of the third cranial nerve, this muscle transitions into an aponeurosis at Whitnall's ligament, inserting into the tarsal plate and skin to create the lid crease.
- **Müller's Muscle:** This sympathetically innervated muscle aids in lid elevation and sits just behind the levator.

The Lacrimal System and Canthal Tendons

The lacrimal gland is situated superolaterally. Surgeons should be wary of a prolapsed gland, which can be mistaken for a lateral fat pad.

The structural integrity of the eye is maintained by the canthal tendons:

- **Medial Canthal Tendon:** A complex, multi-limbed anchor.
- **Lateral Canthal Tendon:** Attaches the tarsal plates to the lateral orbital rim (Whitnall's tubercle). A critical component is the "lateral tarsal strap," which must often be addressed during lower lid tightening or repositioning.




Figure 1.-Lower eye lead anatomy

II. METHODOLOGY

A critical narrative review was conducted using a panel of eight representative scientific articles from the subspecialty of facial plastic and oculoplastic surgery. Inclusion criteria focused on publications addressing: 1) evidence-based assessments of upper blepharoplasty, 2) technical nuances of lower blepharoplasty (including transconjunctival techniques and pinch blepharoplasty), 3) gender considerations in periorbital rejuvenation, and 4) the management of postsurgical complications, specifically eyelid retraction. The information was categorized thematically to facilitate comparative analysis.

III. THEMATIC DEVELOPMENT

Preoperative Evaluation and Risk Reduction Risk mitigation begins with a meticulous preoperative assessment. It is essential to identify preexisting ophthalmological pathologies, paying special attention to dry eye syndrome, which can significantly exacerbate after surgery, especially in patients with a history of refractive surgery (LASIK). In upper blepharoplasty, surgical markings must respect the patient's natural eyelid crease and ethnic characteristics, applying a conservative resection to prevent lagophthalmos and corneal exposure. Furthermore,

Aesthetic Blepharoplasty and Periorbital Rejuvenation: A Comprehensive Review of Surgical Techniques, Risk Prevention, And Complication Management

the surgeon must be prepared to recognize and treat visual emergencies such as retrobulbar hemorrhage.

Upper Blepharoplasty: Preservation and Tarsal Fixation

The goal of modern upper blepharoplasty goes beyond simple age reversal; it seeks to avoid the stigma of the "operated eyelid". Excessive resection of skin and fat interrupts the natural aesthetic continuity. Mendelson and Wong highlight the importance of



Figure 2.-Upper blepharoplasty

stabilizing the eyelid crease (tarsal fixation), which grants independent control over the eyelid segments and avoids reliance on unnatural skin tension. Additionally, the simultaneous approach to volume loss in orbital rim fat or bone support (via augmentation) provides a superior level of aesthetic refinement.

Lower Blepharoplasty: Techniques and Structural Support

The lower eyelid presents unique anatomical challenges due to the effect of gravity and the laxity of canthal support.



CONCLUSIONS

Blepharoplasty has matured from a purely excisional procedure into an intervention of repositioning, preservation, and volumetric support. Current evidence dictates that surgical success depends on limiting musculocutaneous resection and restoring bone and fat contours. Mastery of less invasive approaches, such as the transconjunctival route and the pinch technique, associated with routine canthal suspensions, constitutes the best defence against eyelid malposition's. Continuous education in periorbital anatomy is essential to avoid and, when necessary, manage complex complications such as eyelid retraction.

REFERENCES

- I. Wang Y, Massry GG. Aesthetic Blepharoplasty. *Clin Plast Surg*.
- II. Rao V, Sullivan PK. Lower Lid Blepharoplasty in Men. *Clin Plast Surg*.
- III. Mendelson BC, Wong CH. Upper Blepharoplasty - Nuances for Success. *Clin Plast Surg*.
- IV. Rosenfield LK, Pourriyahi H. The Pinch Blepharoplasty. *Clin Plast Surg*.
- V. Fedok FG. Transconjunctival Lower Blepharoplasty. *Clin Plast Surg*.

Aesthetic Blepharoplasty and Periorbital Rejuvenation: A Comprehensive Review of Surgical Techniques, Risk Prevention, And Complication Management

- VI. Korn BS, Ting M. Reducing Surgical Risks in a Blepharoplasty. *Clin Plast Surg.*
- VII. Steinsapir KD, Steinsapir S. The Treatment of Post-blepharoplasty Lower Eyelid Retraction. *Clin Plast Surg.*
- VIII. Maamari RN. Upper Blepharoplasty: An Evidence-Based Assessment of Preoperative Considerations and Surgical Techniques. *Facial Plast Surg.* 2023;39:273-278